

Walgreens

-PE Product Labels –

-continued

Walgreens

Compare to Maximum Strength
Delsym® Adult Night Time
Cough & Cold active ingredients^{††}

NIGHTTIME NDC 0363-0747-06

Cough & Cold

ACETAMINOPHEN 650 mg /
PAIN RELIEVER, FEVER REDUCER
DIPHENHYDRAMINE HCl 25 mg /
ANTIHISTAMINE, COUGH SUPPRESSANT
PHENYLEPHRINE HCl 10 mg /
NASAL DECONGESTANT

MAXIMUM STRENGTH

- Relieves headaches, fever & sore throat
- Controls cough
- Relieves nasal congestion
- Ages 12 & older

MIXED BERRY FLAVOR

6 FL OZ (177 mL)

PLD-A525A
LB005415

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2018 Walgreen Co.
ITEM 199666 W10013-0418-L



Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey

^{††} This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Delsym® Adult Night Time Cough & Cold.



ORG0418-F

PLD-A525A LB005416

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg.....	Pain reliever/ fever reducer
Diphenhydramine HCl 25 mg.....	Antihistamine/ cough suppressant
Phenylephrine HCl 10 mg.....	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - runny nose
 - sneezing
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other drug containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- new symptoms occur
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for

Drug Facts (continued)

adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product
- dose as follows or as directed by a doctor
- adults and children 12 years and older: 20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

PEEL CORNER FOR MORE DRUG FACTS

WALGREENS NightTime Cough & Cold

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- redness or swelling is present
- new symptoms occur
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with a rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- do not give more than 5 doses in any 24-hour period
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 doses unless directed by a doctor
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- keep dosing cup with product
- dose as follows, or as directed by a doctor
- children 6 to under 12 years of age: 10 mL in dosing cup provided
- children under 6 years of age: do not use

Other information

- each 10 mL contains: sodium 6 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavors, glycerin, propyl galate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING

PARENTS:
Learn about home medicine safety
www.StopMedicineAbuse.org

*Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
**This product is not manufactured or distributed by Reckitt Benckiser, distributor of Children's Mucinex® Cold, Cough & Sore Throat.

NDC 0363-4050-04

Walgreens

Compare to the active ingredients in Children's Mucinex® Cold, Cough & Sore Throat*

Children's Cold, Cough & Sore Throat

ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT
GUAIFENESIN 200 mg / EXPECTORANT
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

- Relief from cough, sore throat, stuffy nose & chest congestion
- Breaks up mucus

AGES 6-11 YEARS

4 FL OZ (118 mL) Berry flavor

NDC 0363-4050-04

Walgreens

Compare to the active ingredients in Children's Mucinex® Cold, Cough & Sore Throat*

Children's Cold, Cough & Sore Throat

ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT
GUAIFENESIN 200 mg / EXPECTORANT
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

- Relief from cough, sore throat, stuffy nose & chest congestion
- Breaks up mucus

AGES 6-11 YEARS

4 FL OZ (118 mL) Berry flavor

Drug Facts

Active Ingredients (in each 10 mL)

Active Ingredients (in each 10 mL)	Purposes
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms
- cough due to minor throat and bronchial irritation
- the intensity of coughing
- the impulse to cough to help your child get to sleep
- nasal congestion
- stuffy nose
- sore throat
- minor aches and pains
- headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has

- liver disease
- heart disease
- high blood pressure
- diabetes
- thyroid disease
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not use more than directed (see Overdose warning).

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.

ITEM 357419 W10024-0322-L



3 11917 16846 3

PLD-E350C F0008008 W3ORG0222-F

Lot No.:
Exp. Date:

COMBO PACK NDC 0363-9015-24



Compare to Maximum Strength Mucinex® Fast-Max® Day Severe Cold & Flu & Maximum Strength Mucinex® Fast-Max® Night Cold & Flu active ingredients!

DAYTIME Cold & Flu	NIGHTTIME Cold & Flu
ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT GUAIFENESIN 200 mg / EXPECTORANT PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT	ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT DOXYLAMINE SUCCINATE 6.25 mg / ANTI-HISTAMINE PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT
<div style="background-color: white; color: #0070C0; padding: 2px; text-align: center; font-weight: bold;">MAXIMUM STRENGTH</div> <ul style="list-style-type: none"> Relieves aches, fever & sore throat Controls cough Thins & loosens mucus Clears nasal & chest congestion 12 years & older <div style="display: flex; align-items: center; margin-top: 20px;"> <div style="text-align: center;"> <div style="font-size: 2em; font-weight: bold; color: white;">16</div> <div style="font-size: 0.8em; color: white;">SOFTGELS</div> </div> <div style="margin: 0 10px;">  </div> <div style="text-align: center;"> <div style="font-size: 0.8em; color: white;">ACTUAL SIZE</div> </div> </div>	<div style="background-color: white; color: #002060; padding: 2px; text-align: center; font-weight: bold;">MAXIMUM STRENGTH</div> <ul style="list-style-type: none"> Relieves aches, fever, sore throat, nasal congestion, runny nose & sneezing Controls cough 12 years & older <div style="display: flex; align-items: center; margin-top: 20px;"> <div style="text-align: center;"> <div style="font-size: 2em; font-weight: bold; color: white;">8</div> <div style="font-size: 0.8em; color: white;">SOFTGELS</div> </div> <div style="margin: 0 10px;">  </div> <div style="text-align: center;"> <div style="font-size: 0.8em; color: white;">ACTUAL SIZE</div> </div> </div>
<div style="font-size: 1.5em; font-weight: bold;">TOTAL 24</div> <div style="font-size: 0.8em;">SOFTGELS</div>	

3227

Walgreens

DAYTIME
Cold & Flu

150501005400

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleepiness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 softgels in any 24-hour period
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients FD&C yellow #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide

Questions or Comments? Call 1-877-296-4088

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2022 Walgreen Co.
MADE IN INDIA

6907 0322

W3ORG0522-F

Walgreens

COMBO PACK

NDC 0383-0775-24

DAYTIME
Cold & Flu

NIGHTTIME
Cold & Flu

1006109

Drug Facts

Active ingredients (in each softgel) Purposes

Acetaminophen 325 mg... Pain reliever/fever reducer
Dextromethorphan HBr 10 mg... Cough suppressant
Doxylamine succinate 6.25 mg... Antihistamine
Phenylephrine HCl 5 mg... Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - sinus congestion and pressure
 - cough
 - minor aches and pains
 - headache
 - nasal congestion
 - sore throat
 - runny nose
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- controls cough to help you get to sleep
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 12 softgels in 24 hours, which is the maximum daily amount.

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to enlarged prostate gland
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Maximum Strength

- Relieves aches, fever, & sore throat
- Controls cough
- Thins & loosens mucus
- Clears nasal & chest congestion
- 12 years & older

Maximum Strength

- Relieves aches, fever, sore throat, nasal congestion, runny nose & sneezing
- Controls cough
- 12 years & older

16 SOFTGELS* (LIQUID-FILLED CAPSULES)

8 SOFTGELS* (LIQUID-FILLED CAPSULES)

TOTAL 24 SOFTGELS

ITEM 428154 W00000-0000-0

LOT NO.:

EXP DATE:

Walgreens

NIGHTTIME
Cold & Flu

150501005400

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleepiness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 softgels in any 24-hour period
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide

Questions or Comments? Call 1-877-296-4088

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2022 Walgreen Co.
MADE IN INDIA

6907 0322

W3ORG0522-F

COMBO PACK NDC 0363-9014-24

Walgreens

Compare to Maximum Strength
Mucinex® Sinus-Max® Day & Mucinex®
Sinus-Max® Night active ingredients††

DAYTIME

Sinus Relief

ACETAMINOPHEN 325 mg /
PAIN RELIEVER
DEXTROMETHORPHAN HBr 10 mg /
COUGH SUPPRESSANT
GUAIFENESIN 200 mg /
EXPECTORANT
PHENYLEPHRINE HCl 5 mg /
NASAL DECONGESTANT

NIGHTTIME

Sinus Relief

ACETAMINOPHEN 325 mg /
PAIN RELIEVER
DEXTROMETHORPHAN HBr 10 mg /
COUGH SUPPRESSANT
DOXYLAMINE SUCCINATE 6.25 mg /
ANTIHISTAMINE
PHENYLEPHRINE HCl 5 mg /
NASAL DECONGESTANT

MAXIMUM STRENGTH

- Relieves sinus pressure, headache & congestion
- Controls cough
- Thins & loosens mucus
- 12 years & older

16
SOFTGELS


ACTUAL SIZE

MAXIMUM STRENGTH

- Relieves nasal congestion, sinus pressure & pain, runny nose & sneezing
- Controls cough
- 12 years & older

8
SOFTGELS


ACTUAL SIZE

TOTAL 24 SOFTGELS

Walgreens Cold & Congestion Relief Drug Facts

Drug Facts (continued)

Directions

- do not take more than 6 doses in any 24 hour period
- do not exceed recommended dosage
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- 1 mL = milliliter

Age	Dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

Other information

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F)
- do not refrigerate
- protect from light

Inactive ingredients acacia, caffeine, potassium, anhydrous citric acid, EDTA disodium, FD&C blue #1, FD&C red #40, flavor, methylcellulose, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium chloride, xanthan gum

Questions or comments?
Call 1-877-753-3836 Monday-Friday 9AM-5PM EST

"This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Children's Dimetane® Cold & Cough."

Walgreens Cold & Cough Relief Drug Facts

Drug Facts

Active ingredients (in each 10 mL)
Brompheniramine maleate 2 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purposes
Antihistamine
Cough suppressant
Nasal decongestant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis)
- runny nose
- sneezing
- itching of the nose or throat
- temporarily restores free breathing through the nose

Warnings
Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking any other oral nasal decongestant or stimulant
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

Walgreens Cold & Cough Relief Drug Facts

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- 1 mL = milliliter

Age	Dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to 11 years	10 mL every 4 hours
children under 6 years	do not use

Other information

- each 10 mL contains: sodium 5 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose

Questions or comments?
Call 1-877-753-3836 Monday-Friday 9AM-5PM EST

TAMPER EVIDENT: DO NOT USE IF CAPTION IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

"Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands."
"This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Children's Dimetane® Nighttime Cold & Congestion."

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
©2021 Walgreen Co.

DAY & NIGHT PACK NDC 0363-4500-08

Walgreens 

Compare to the active ingredients in Children's Dimetane® Cold & Nighttime Cold & Congestion®

Children's Cold & Cough Relief
BROMPHENIRAMINE MALEATE 2 mg / ANTIHISTAMINE
DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

Children's Cold & Congestion Relief
DIPHENHYDRAMINE HCl 12.5 mg / ANTIHISTAMINE / COUGH SUPPRESSANT
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

Daytime
Alcohol Free

- Relieves nasal congestion, runny nose, itchy & watery eyes, sneezing & coughing

Nighttime
Alcohol Free

- Relieves stuffy nose, runny nose, sneezing, itchy & watery eyes & coughing

AGES 6 YEARS & OLDER Grape flavor

AGES 6 YEARS & OLDER Grape flavor

2 - 4 FL OZ (118 mL) BOTTLES / TOTAL 8 FL OZ (236 mL)

ITEM 904373 W10024-0322-L



3 11917 17679 6



PAPER PLASTIC PLASTIC CUP

Not recycled in all communities

PLD-E2610
PLD-E2610A
PC000013

Lot No.:
Exp. Date:



Drug Facts

Active Ingredients (In each softgel)

- Acetaminophen 325 mg
- Dextromethorphan HBr 10 mg
- Guaifenesin 200 mg
- Phenylephrine HCl 5 mg

Purpose

- Pain reliever/fever reducer
- Cough suppressant
- Expectorant
- Nasal decongestant



Drug Facts (continued)

Uses temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- minor aches & pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores free breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to help the bronchial passageways of both nose and mouth make coughs more productive.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 softgels in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- you get dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 8 softgels per 24 hours

adults & children 12 years & over	2 softgels with water every 4 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

When using other Nighttime or Daytime products, carefully read each label to ensure correct dosing.

Other information do not exceed 25°C

Inactive ingredients

FD&C blue #1, FD&C red #40, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide

Questions or Comments?

Call 1-877-290-4008



Print/Varnish Omit Area

THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT AND TAMPER-EVIDENT PACKAGE. USE ONLY IF BLISTERS ARE INTACT.

Varnish Omit Area

NDC 0363-0109-24

Walgreens

Compare to the active ingredients in Vicks® DayQuil® Severe Cold & Flu†

DAYTIME Severe Cold & Flu

ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER
 DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT
 GUAIFENESIN 200 mg / EXPECTORANT
 PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

Maximum Strength

- Relieves headache, fever, sore throat, minor aches & pains, cough, chest congestion, thin & loosens mucus, nasal congestion & sinus pressure

24 SOFTGELS

ACTUAL SIZE

W3ORG0522-F

Varnish Omit Area

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING

This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex®
Fast-Max® Severe Cold and Night Time Cold & Flu.

3232

Nighttime Cold & Flu (continued)		Daytime Severe Cold		Daytime Severe Cold (continued)		DAY & NIGHT PACK		Nighttime Cold & Flu	
Drug Facts (continued) When using this product ■ do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. Overdose warnings: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Directions ■ do not take more than directed (see Overdose warning) ■ do not take more than 12 softgels (Daytime and Nighttime) in any 24-hour period ■ adults and children 12 years of age and older: take 2 softgels every 4 hours ■ children under 12 years of age: do not use ■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing Other information ■ swallow whole; do not crush, chew, or dissolve ■ store between 15-30°C (59-86°F) ■ avoid excessive heat Inactive ingredients DAC yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol, sorbitol, titanium dioxide *contains one or more of these ingredients Questions or comments? Call 1-877-753-3838 Monday-Friday 9AM-5PM EST TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING. KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION		Drug Facts Active ingredients (in each softgel) Acetaminophen 325 mg. Pain reliever/fever reducer Dextromethorphan HBr 10 mg. Cough suppressant Pseudoephedrine HCl 5 mg. Nasal decongestant Purposes ■ temporarily relieves these common cold and flu symptoms ■ headache ■ nasal congestion ■ sore throat ■ minor aches and pains ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive ■ temporarily reduces fever Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have ■ liver disease ■ diabetes ■ high blood pressure ■ heart disease ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ a persistent or chronic		Drug Facts (continued) When using this product ■ do not use more than directed. Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Directions ■ do not take more than directed (see Overdose warning) ■ do not take more than 12 softgels (Daytime and Nighttime) in any 24-hour period ■ adults and children 12 years of age and older: take 2 softgels every 4 hours ■ children under 12 years of age: do not use ■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing Other information ■ swallow whole; do not crush, chew, or dissolve ■ store between 15-30°C (59-86°F) ■ avoid excessive heat Inactive ingredients FD&C red #40, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, sorbitol, titanium dioxide *may contain this ingredient Questions or comments? Call 1-877-753-3838 Monday-Friday 9AM-5PM EST Walgreens Pharmacist Recommended Walgreens Pharmacist Survey This product is not manufactured or distributed by Rectivix Biotech, distributor of Maximum Strength Mucinex® Fast-Max® Day Severe Cold and Maximum Strength Mucinex® Fast-Max® Night Cold & Flu.		Walgreens Compare to Maximum Strength Mucinex® Fast-Max® Day Severe Cold & Maximum Strength Mucinex® Fast-Max® Night Cold & Flu active ingredients! DAYTIME Severe Cold ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT GUAFENESIN / EXPECTORANT PHENYLEPHRINE HCl / NASAL DECONGESTANT MAXIMUM STRENGTH ALCOHOL FREE • Relieves aches, fever & sore throat • Controls cough • Clears nasal & chest congestion • Thins & loosens mucus • Ages 12 & over 16 SOFTGELS* (LIQUID-FILLED CAPSULES) ACTUAL SIZE NIGHTTIME Cold & Flu ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT DOXYLAMINE SUCCIATE / ANTIHISTAMINE PHENYLEPHRINE HCl / NASAL DECONGESTANT MAXIMUM STRENGTH ALCOHOL FREE • Relieves aches, fever, sore throat, nasal congestion, runny nose & sneezing • Controls cough • Ages 12 & over 8 SOFTGELS* (LIQUID-FILLED CAPSULES) ACTUAL SIZE 16 SOFTGELS* - 8 SOFTGELS* / TOTAL 24 SOFTGELS*		Drug Facts Active ingredients (in each softgel) Acetaminophen 325 mg. Pain reliever/fever reducer Dextromethorphan HBr 10 mg. Cough suppressant Doxylamine succinate 6.25 mg. Antihistamine Phenylephrine HCl 5 mg. Nasal decongestant Purposes ■ temporarily relieves these common cold and flu symptoms ■ cough ■ headache ■ minor aches and pains ■ sore throat ■ nasal congestion ■ runny nose and sneezing ■ controls cough to help you get to sleep ■ temporarily reduces fever Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have ■ liver disease ■ diabetes ■ high blood pressure ■ heart disease ■ glaucoma ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough that occurs with too much phlegm (mucus) Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers	

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
Walgreens
100% SATISFACTION GUARANTEED
walgreens.com ©2018 Walgreen Co.

ITEM 946682 W10024-1018-L
3 11917 18621 4

PLD-0421C

FD006388

Lot No.:

Exp. Date:



Nighttime Multi-Symptom Cold Drug Facts

Drug Facts (continued)

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- neurosis, dizziness, or sleeplessness occur
- stomach, heart, or lung problems occur
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- do not give more than 5 doses in any 24-hour period
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 doses unless directed by a doctor
- measure only with dosing cup provided. Do not use any other dosing device.
- ml, = milliliter
- do as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL every 4 hours
- children under 6 years of age: do not use

Other information

- each 10 mL contains sodium 6 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucrose, xanthan gum

Daytime Multi-Symptom Cold Drug Facts

Drug Facts

Active ingredients (in each 5 mL) Purposes

Dextromethorphan HBr 5 mg Cough suppressant

Guaifenesin 100 mg Expectorant

Phenylephrine HCl 2.5 mg Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of thickened mucus and make coughs more productive
- temporarily relieves
- cough due to minor throat and bronchial irritation as may occur with the common cold or irritated infants
- the intensity of coughing
- the impulse to cough to help your child get to sleep
- nasal congestion due to a cold
- stuffy nose

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if your child has

- heart disease
- high blood pressure
- diabetes
- thyroid disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product, do not use more than directed.

Stop use and ask a doctor if

- neurosis, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with a fever, rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose: get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not use more than 6 doses in any 24-hour period
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- keep dosing cup with product
- do as follows or as directed by a doctor
- ml, = milliliter
- Age
- children 6 years to under 12 years 10 mL every 4 hours
- children 4 years to under 6 years 5 mL every 4 hours
- children under 4 years do not use

Daytime Multi-Symptom Cold Drug Facts

Drug Facts (continued)

Other information

- each 5 mL contains: sodium 5 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, disodium EDTA, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucrose, xanthan gum

Questions or comments?
Call 1-877-753-3086 Monday-Friday 9AM-5PM EST

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

KEEP OUT OF REACH OF CHILDREN. READ COMPLETE WARNINGS AND PRODUCT INFORMATION.

DOSAGE CUP PROVIDED

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by Rinkia Benckiser, distributor of Children's Mucinex® Night Time Multi-Symptom Cold and Children's Mucinex® Multi-Symptom Cold.

DISTRIBUTED BY: WALGREEN CO., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED

walgreens.com

©2021 Walgreen Co.

DAY & NIGHT PACK

NDC 0363-4200-08

Walgreens

Compare to the active ingredients in Children's Mucinex® Multi-Symptom Cold & Night Time Multi-Symptom Cold

MULTI-SYMPTOM Children's Cold

DEXTROMETHORPHAN HBr 5 mg / COUGH SUPPRESSANT
GUAFENESIN 100 mg / EXPECTORANT
PHENYLEPHRINE HCl 2.5 mg / NASAL DECONGESTANT

Daytime

- Alcohol Free
- Relief from chest congestion, cough & stuffy nose
- Breaks up mucus

MULTI-SYMPTOM Children's Cold

ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER
DIPHENHYDRAMINE HCl 12.5 mg / ANTIHISTAMINE / COUGH SUPPRESSANT
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

Nighttime

- Alcohol Free
- Relief from stuffy nose, cough, fever, sore throat, runny nose & sneezing

AGES 4 - 11 YEARS Berry flavor

AGES 6 - 11 YEARS Berry flavor

2 - 4 FL OZ (118 mL) BOTTLES / TOTAL 8 FL OZ (236 mL)

Nighttime Multi-Symptom Cold Drug Facts

Drug Facts

Active ingredients (in each 10 mL) Purposes

Acetaminophen 325 mg Pain reliever/fever reducer

Diphenhydramine HCl 12.5 mg Antihistamine/Cough suppressant

Phenylephrine HCl 5 mg Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
- minor aches and pains
- headache
- sore throat
- sneezing
- runny nose
- nasal congestion
- cough
- controls cough to help your child get to sleep
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- diarrhea
- itching
- if a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- with any other drug containing diphenhydramine even one used on the skin
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product
- to make a child sleepy

Ask a doctor before use if your child has

- liver disease
- heart disease
- high blood pressure
- diabetes
- thyroid disease
- glaucoma
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if your child is

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

<p>▲ When using this product, do not use more than directed.</p> <p>▲ Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p> <p>▲ Chronic bronchitis or emphysema</p>	<p>▲ exhaustion may occur, especially in children</p> <p>▲ marked drowsiness may occur</p> <p>▲ avoid alcoholic drinks</p> <p>▲ be careful when driving a motor vehicle or operating machinery</p> <p>▲ alcohol, sedatives, and tranquilizers may increase drowsiness</p>
---	--

MSD1161522-F

- ASCENDING 1515

KEEP OUTER CARTON FOR COMPLETE
WARNINGS AND PRODUCT INFORMATION



Nighttime Sinus Relief (continued)

Drug Facts (continued)

Stop use and ask a doctor if: ■ nervousness, dizziness, or sleeplessness occur ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed (see Overdose warning) ■ do not take more than 12 softgels (Daytime and Nighttime) in any 24-hour period ■ adults and children 12 years of age and older: take 2 softgels every 4 hours ■ children under 12 years of age: do not use ■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information

■ swallow whole; do not crush, chew, or dissolve ■ store between 15-30°C (59-86°F) ■ avoid excessive heat

Inactive ingredients

D&C yellow #10, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol, sorbitol, titanium dioxide

Questions or comments?

Call 1-877-753-3835 Monday-Friday 9AM-5PM EST

Pharmacist Recommended

Health expertise you rely on, quality you trust.

Daytime Sinus Relief

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg	Pain reliever
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

■ temporarily relieves ■ nasal congestion ■ headache ■ cough due to inhaled irritants ■ sinus congestion and pressure ■ minor aches and pains ■ promotes nasal and/or sinus drainage ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

†Walgreens Pharmacist Survey
†This product is not manufactured or distributed by Walgreens. Walgreens is the distributor of Maximum Strength Mucinex® Sinus-Max® Day and Maximum Strength Mucinex® Sinus-Max® Night.

Daytime Sinus Relief (continued)

Drug Facts (continued)

Ask a doctor before use if you have: ■ liver disease ■ high blood pressure ■ diabetes ■ heart disease ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if: ■ nervousness, dizziness, or sleeplessness occur ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed (see Overdose warning) ■ do not take more than 12 softgels (Daytime and Nighttime) in any 24-hour period ■ adults and children 12 years of age and older: take 2 softgels every 4 hours ■ children under 12 years of age: do not use ■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information

■ swallow whole; do not crush, chew, or dissolve ■ store between 15-30°C (59-86°F) ■ avoid excessive heat

Inactive ingredients

FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, sorbitol, titanium dioxide

Questions or comments?

Call 1-877-753-3835 Monday-Friday 9AM-5PM EST

DAY & NIGHT PACK

NDC 0363-0723-24

Walgreens

Compare to Maximum Strength Mucinex® Sinus-Max® Day & Mucinex® Sinus-Max® Night active ingredients!

DAYTIME Sinus Relief

ACETAMINOPHEN / PAIN RELIEVER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
GUAIFENESIN / EXPECTORANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

MAXIMUM STRENGTH
ALCOHOL FREE

- Relieves sinus pressure, headache & congestion
- Controls cough
- Thins & loosens mucus
- Ages 12 & over

16 SOFTGELS* ACTUAL SIZE

NIGHTTIME Sinus Relief

ACETAMINOPHEN / PAIN RELIEVER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
DOXYLAMINE SUCCINATE / ANTIHISTAMINE
PHENYLEPHRINE HCl / NASAL DECONGESTANT

MAXIMUM STRENGTH
ALCOHOL FREE

- Relieves nasal congestion, sinus pressure & pain, runny nose & sneezing
- Controls cough
- Ages 12 & over

8 SOFTGELS* ACTUAL SIZE

TOTAL 24 SOFTGELS* (*LIQUID-FILLED CAPSULES)

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
©2017 Walgreen Co.

ITEM 946681 W10024-0118-L

3 11917 18620 7

PARENTS:
www.3softgelsdayandnight.com

PLD-C425B
FC004407

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Lot No.:
Exp. Date:

Nighttime Sinus Relief

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg	Pain reliever
Dextromethorphan HBr 10 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

■ temporarily relieves ■ nasal congestion ■ cough due to inhaled irritants ■ headache ■ sinus congestion and pressure ■ minor aches and pains ■ runny nose and sneezing ■ promotes nasal and/or sinus drainage

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have: ■ liver disease ■ high blood pressure ■ diabetes ■ heart disease ■ thyroid disease ■ glaucoma ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are: ■ taking sedatives or tranquilizers

When using this product: ■ do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery

Walgreens

DAYTIME Sinus Relief

NIGHTTIME Sinus Relief

COMBO PACK

Walgreens

Compare to the active ingredients in Maximum Strength Brand's® Sinus-Max® Day & Nighttime Sinus-Max® Night

DAYTIME Sinus Relief

NIGHTTIME Sinus Relief

ACETAMINOPHEN 325 mg / PAIN RELIEVER

DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT

GUAFENESIN 200 mg / EXPECTORANT

PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

Directions

Do not take more than directed (see Overdose Warning)

Other information

Inactive ingredients

Questions or Comments? Call 1-877-296-4008

DISTRIBUTED BY WALGREENS CO., 200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED

www.walgreens.com ©2022 Walgreen Co. MADE IN INDIA

WSORG0522-F

Drug Facts

Active ingredients (in each softgel)

Uses

Warnings

Do not use

Ask a doctor before use if you have

Ask a doctor or pharmacist before use if you are

Other information

Inactive ingredients

Questions or Comments? Call 1-877-296-4008

Parents: Learn about teen medicine abuse

Parents: Learn about teen medicine abuse

Parents: Learn about teen medicine abuse

Over-the-counter convenience:
Call 1-877-763-3435
Distributors: WALGREENS CO.,
200 W. WATSON ST., DEERFIELD, IL 60015
www.walgreens.com ©2011 Walgreen Co.
ITEM# 007515 NOV000000-003

DAY & NIGHT PACK

Walgreens

Compare to the active ingredients in Maximum Strength Mucinex® Fast-Act® Severe Congestion & Cough & Hixmoxin® Cold & Flu*

DAYTIME

Severe Congestion & Cough

DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
GUAFENESIN / EXPECTORANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

Maximum Strength

- Controls cough
- Clears nasal & chest congestion
- Thins & loosens mucus
- 12 years & older

NIGHTTIME

Cold & Flu

ACETAMINOPHEN / PAIN RELIEVER & FEVER REDUCER
DIPHENHYDRAMINE HCl / ANTIHISTAMINE / COUGH SUPPRESSANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

Maximum Strength

- Relieves aches, fever, sore throat, nasal congestion, runny nose & sneezing
- Controls cough
- 12 years & older

2 - 6 FL OZ (177 mL) BOTTLES / TOTAL 12 FL OZ (354 mL)

*Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
*This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademarks Mucinex® and Fast-Max®.

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Day Facts (continued) Ingredients: See the inactive ingredients on the back of this bottle.

- **Directions:** • **Use:** • **Warnings:** • **Precautions:** • **Interactions:** • **Side Effects:** • **Storage:** • **Expiration:** • **Lot:** • **Manufactured by:** • **Walgreen Company**

Walgreens

Compare to Maximum Strength
Mucinex® Fast-Max® Cold & Flu
ALL IN ONE active ingredients††

NDC 0363-9017-16

Cold & Flu

NEW

ACETAMINOPHEN / PAIN RELIEVER & FEVER REDUCER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
GUAIFENESIN / EXPECTORANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

MULTI-SYMPTOM

MAXIMUM STRENGTH

DISSOLVES QUICKLY

- Relieves: Headache, body pain, fever, sore throat, chest congestion, cough, nasal congestion, sinus congestion & sinus pressure
- 12 years & older

16 SOFTGELS



ACTUAL SIZE

Walgreens

*Compare to Maximum Strength Mucinex® FAST-MAX®
Congestion & Headache Liquid Gels active ingredients!†*

NDC 0363-9019-16

NEW

Congestion & Headache

ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

MAXIMUM STRENGTH

- Relieves headache & fever
- Controls cough
- Relieves nasal congestion
- 12 years & older

16 SOFTGELS*
(*LIQUID-FILLED CAPSULES)



ACTUAL SIZE

3244

NDC 0363-0512-06

Drug Facts**Active ingredients (in each 5 mL)**

Dextromethorphan HBr 5 mg.....Cough suppressant
Guaifenesin 100 mg.....Expectorant
Phenylephrine HCl 2.5 mg.....Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help your child get to sleep
 - nasal congestion due to a cold ■ stuffy nose

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if your child has

- heart disease ■ high blood pressure ■ thyroid disease ■ diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product, do not use more than directed.

Stop use and ask a doctor if

- your child gets nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with a fever, rash, or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 6 doses in any 24-hour period
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- keep dosing cup with product
- dose as follows or as directed by a doctor ■ mL = milliliter

Age	Dose
children 6 years to under 12 years	10 mL every 4 hours
children 4 years to under 6 years	5 mL every 4 hours
children under 4 years	do not use

Other information

- each 5 mL contains: sodium 5 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, disodium EDTA, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call 1-877-753-3835 Monday-Friday 9AM-5PM EST

¹Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

²This product is not manufactured or distributed by Reckitt Benckiser, distributor of Children's Mucinex® Congestion & Cough.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION



DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED

walgreens.com

©2021 Walgreen Co.

Walgreens



Compare to the active ingredients in
Children's Mucinex® Congestion & Cough²

Children's Mucus Congestion & Cough Relief

DEXTROMETHORPHAN HBr 5 mg /
COUGH SUPPRESSANT
GUAIFENESIN 100 mg / EXPECTORANT
PHENYLEPHRINE HCl 2.5 mg /
NASAL DECONGESTANT

Alcohol Free

- Relief from chest congestion, cough & stuffy nose
- Breaks up mucus

AGES
4-11
YEARS



6.8 FL OZ (201 mL)



DOSAGE CUP
PROVIDED

PARENTS:

Learn about teen medicine abuse
www.StopMedicineAbuse.org

ITEM 593323 W10024-0322-L



PLD-BS45A FC008012

Lot No.:

Exp. Date:

[illegible]

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING

3247

Walgreens

Compare to Maximum Strength Mucinex® Fast-Max® All-In-One Cold & Flu active ingredients!

DAYTIME NDC 0363-0886-06

Cold & Flu

ACETAMINOPHEN 650 mg / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr 20 mg / COUGH SUPPRESSANT
GUAIFENESIN 400 mg / EXPECTORANT
PHENYLEPHRINE HCl 10 mg / NASAL DECONGESTANT

MAXIMUM STRENGTH

- Relieves aches, fever, sore throat, nasal & chest congestion, sinus congestion & pressure
- Controls cough
- Thins & loosens mucus
- 12 years & older

6 FL OZ (177 mL) LIQUID

†† This product is not manufactured or distributed by Reckitt Benckiser, distributor of Mucinex® Maximum Strength Mucinex® Fast-Max® All-In-One Cold & Flu. PLD-A574A L8006719 ITEM 907517 W10389-10/15-F ORG1019-F

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Mucinex® Fast-Max® Maximum Strength All-In-One Cold & Flu.

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

PLD-A574A L8006719

Drug Facts**Active ingredients (in each 20 mL)**

Acetaminophen 650 mg.....Pain reliever/fever reducer
 Dextromethorphan HBr 20 mg.....Cough suppressant
 Guaifenesin 400 mg.....Expectorant
 Phenylephrine HCl 10 mg.....Nasal decongestant

Purposes

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)**Uses**

- temporarily relieves these common cold and flu symptoms
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - stuffy nose
 - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

When using this product, do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period

Drug Facts (continued)

- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

3248

Walgreens

Compare to Maximum Strength
Mucinex® Fast-Max® All-In-One Night
Time Cold & Flu active ingredients††

NIGHTTIME NDC 0363-9130-06

Cold & Flu

ACETAMINOPHEN 650 mg /
PAIN RELIEVER / FEVER REDUCER
DIPHENHYDRAMINE HCl 25 mg /
ANTIHISTAMINE / COUGH SUPPRESSANT
PHENYLEPHRINE HCl 10 mg /
NASAL DECONGESTANT

MAXIMUM STRENGTH

- Relieves cough, fever, sore throat, body pain, sneezing, itchy throat, headache, nasal congestion & runny nose
- 12 years & older

6 FL OZ (177 mL) LIQUID

†† This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® All-In-One Night Time Cold & Flu. PLD-A603A LB006368 ITEM 907517 W10369-10/15-F ORG1019-F

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Night Time All In One Cold & Flu.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

PLD-A603A LB006360

Drug Facts

Active ingredients (in each 20 mL)

Acetaminophen 650 mg.....Pain reliever/fever reducer
Diphenhydramine HCl 25 mg.....Antihistamine/
cough suppressant
Phenylephrine HCl 10 mg.....Nasal decongestant

Purposes

PEEL CORNER FOR MORE DRUG FACTS ▶

Drug Facts (continued)

Uses

- temporarily relieves these common cold and flu symptoms
- cough ■ nasal congestion
- minor aches and pains ■ sore throat
- headache ■ runny nose ■ sneezing
- sinus congestion and pressure
- itching of the nose or throat
- itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away. ▶

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other drug containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease ■ heart disease ■ diabetes
- thyroid disease ■ high blood pressure
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus) ▶

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or ▶

Drug Facts (continued)

contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product ■ mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years and older:
20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

PEEL CORNER FOR MORE DRUG FACTS ▶

NDC 0363-3361-06



Compare to the active ingredients in
Maximum Strength Mucinex® Fast-Max®
Cold, Flu & Sore Throat¹

MUCUS RELIEF

Cold, Flu & Sore Throat

**ACETAMINOPHEN 650 mg /
PAIN RELIEVER & FEVER REDUCER
DEXTROMETHORPHAN HBr 20 mg /
COUGH SUPPRESSANT / GUAIFENESIN 400 mg /
EXPECTORANT / PHENYLEPHRINE HCl 10 mg /
NASAL DECONGESTANT**

Maximum Strength

- Relieves headache & fever
- Controls cough
- Relieves nasal & chest congestion
- Thins & loosens mucus
- For ages 12 & over

6 FL OZ (177 mL)

PLD-B282C
LB008816

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

©2022 Walgreen Co.

¹Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

²This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademarks Mucinex® and Fast-Max®.

Discard Seal,
Empty &
Replace Cap



PLASTIC BOTTLE
PLASTIC CUP

how2recycle.info LB008817 PLD-B282C



Drug Facts

Active ingredients (in each 20 mL)

Acetaminophen 650 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....Cough suppressant
Guaifenesin 400 mg.....Expectorant
Phenylephrine HCl 10 mg.....Nasal decongestant

Purposes

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

Uses

- temporarily relieves these common cold and flu symptoms
- sinus congestion and pressure
- minor aches and pains
- nasal congestion
- cough due to minor throat and bronchial irritation
- sore throat
- headache
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Drug Facts (continued)

When using this product, do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period

Drug Facts (continued)

- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

PEEL CORNER FOR MORE DRUG FACTS

Walgreens

Compare to Maximum Strength Mucinex®
Fast-Max® Clear & Cool® Cold,
Flu & Sore Throat active ingredients†

NDC 0083-0782-06

Cold, Flu & Sore Throat

ACETAMINOPHEN 650 mg /
PAIN RELIEVER & FEVER REDUCER
DEXTROMETHORPHAN HBr 20 mg /
COUGH SUPPRESSANT
GUAIFENESIN 400 mg / EXPECTORANT
PHENYLEPHRINE HCl 10 mg / NASAL DECONGESTANT

MAXIMUM STRENGTH

- Relieves headache & fever
- Controls cough
- Relieves nasal & chest congestion
- Thins & loosens mucus
- 12 years & older

MINT FLAVOR

6 FL OZ (177 mL)

PLD-C452B
LB006897

3250

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY
SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2019 Walgreen Co.
ITEM 984979 W10139-0719-F



Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey

†This product is not manufactured or
distributed by Reckitt Benckiser,
distributor of Maximum Strength
Mucinex® Fast-Max® Clear & Cool®
Cold, Flu & Sore Throat.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

ORG0719-F

PLD-C452B LB006896

PEEL CORNER FOR DRUG FACTS

Drug Facts**Active ingredients
(in each 20 mL)****Purposes**

Acetaminophen 650 mg.....	Pain reliever/ fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms
 - sinus congestion and pressure
 - minor aches and pains
 - headache
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

Drug Facts (continued)

- more than 4,000 mg of acetaminophen in 24 hours
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks daily while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure

Drug Facts (continued)

- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are
taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for

Drug Facts (continued)

adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, disodium EDTA, FD&C blue #1, FD&C yellow #10, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

PEEL CORNER FOR MORE DRUG FACTS

NDC 0363-0337-06

Walgreens

Compare to the active ingredients in
Maximum Strength Mucinex® Fast-Max®
Severe Congestion & Cough[†]

**Severe
Congestion
& Cough**

DEXTROMETHORPHAN HBr /
COUGH SUPPRESSANT
GUAIFENESIN / EXPECTORANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

Maximum Strength

- Controls cough
- Clears nasal & chest congestion
- Thins & loosens mucus
- 12 years & older

6 FL OZ (177 mL)

PLD-E283D
LB008701

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY
SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.
ITEM 280506 W10139-0122-F



Walgreens Pharmacist Recommended.
Our pharmacists recommend the
Walgreens brand. We invite you to
compare to national brands.
^{††}This product is not manufactured or
distributed by RB Health (US) LLC, owner of
the registered trademarks Mucinex® and
Fast-Max®.

Discard Seals,
Empty &
Replace Cap



PLASTIC
BOTTLE



PLASTIC
CUP

how2recycle.info

PARENTS:

Learn about teen medicine abuse
www.StopMedicineAbuse.org

W3ORG0122-F

PLD-E283D LB008703

PEEL CORNER FOR DRUG FACTS

Drug Facts

Active ingredients Purposes (in each 20 mL)

Dextromethorphan HBr 20 mg.....
.....Cough suppressant
Guaifenesin 400 mg.....Expectorant
Phenylephrine HCl 10 mg.....
.....Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin
bronchial secretions to rid the bronchial
passageways of bothersome mucus
and make coughs more productive
- temporarily relieves
 - cough due to minor throat and
bronchial irritation as may occur
with the common cold or inhaled
irritants
 - the intensity of coughing
 - the impulse to cough to help you
get to sleep
 - nasal congestion due to a cold

Drug Facts (continued)

Warnings

Do not use

- if you are now taking a prescription
monoamine oxidase inhibitor (MAOI)
(certain drugs for depression,
psychiatric or emotional conditions, or
Parkinson's disease), or for 2 weeks
after stopping the MAOI drug. If you do
not know if your prescription drug
contains an MAOI, ask a doctor or
pharmacist before taking this product.
- for children under 12 years of age

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged
prostate gland
- persistent or chronic cough such as
occurs with smoking, asthma, chronic
bronchitis, or emphysema
- cough that occurs with too much
phlegm (mucus)

When using this product, do not use
more than directed.

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or
sleeplessness occur
- symptoms do not get better within
7 days or occur with fever
- cough comes back, or occurs with
fever, rash, or headache that lasts.
These could be signs of a serious
condition.

If pregnant or breast-feeding, ask a
health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or
contact a Poison Control Center
(1-800-222-1222) right away.

Directions

- do not take more
than 6 doses in any 24-hour period
- measure only with dosing cup provided.
Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- shake well before using
- adults and children 12 years of age and
older: 20 mL in dosing cup provided
every 4 hours
- children under 12 years of age:
do not use

Drug Facts (continued)

Other information

- each 20 mL contains: sodium 17 mg
- store between 20-25°C (68-77°F). Do
not refrigerate.

Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1,
FD&C red #40, flavor, glycerin, propyl
gallate, propylene glycol, purified water,
sodium benzoate, sodium citrate, sorbitol,
sucralose, xanthan gum

Questions or comments?

Call 1-877-753-3935 Monday-Friday
9AM-5PM EST

PEEL CORNER FOR MORE DRUG FACTS

NDC 0363-0681-06

Walgreens

Compare to the active ingredients in
Maximum Strength Mucinex® Fast-Max®
Clear & Cool® Severe Congestion & Cough®

Severe Congestion & Cough

DEXTROMETHORPHAN HBr 20 mg /
COUGH SUPPRESSANT
GUAIFENESIN 400 mg / EXPECTORANT
PHENYLEPHRINE HCl 10 mg /
NASAL DECONGESTANT

Maximum Strength

- Thins & loosens mucus
- Relieves nasal & chest congestion
- Controls cough
- 12 years & older

6 FL OZ (177 mL) Mint flavor

PLD-D453A LB008711

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.
ITEM 984978 W10139-0222-F



Walgreens Pharmacist Recommended.
Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

*This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademarks Mucinex®, Fast-Max® and Clear & Cool®.

Discard Seal,
Empty &
Replace Cap



PLASTIC
BOTTLE

*Not recycled in
all communities



PLASTIC
CUP

*Not recycled in
all communities

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

W3ORG0122-F

PLD-D453A LB008710

PEEL CORNER FOR DRUG FACTS

Drug Facts

Active ingredients (in each 20 mL)

Dextromethorphan HBr 20 mg.....Cough suppressant
Guaifenesin 400 mg.....Expectorant
Phenylephrine HCl 10 mg.....Nasal decongestant

Purposes

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - nasal congestion due to a cold
 - temporarily helps you cough less

Drug Facts (continued)

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product, do not use more than directed.

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Drug Facts (continued)

Other information

- each 20 mL contains: sodium 18 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, disodium EDTA, FD&C blue #1, FD&C yellow #10, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

PEEL CORNER FOR MORE DRUG FACTS



TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL ABOVE OR UNDER CAP IS BROKEN OR MISSING

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.
ITEM 907515 W10139-0222-F



W3ORG0122-F PLD-E377D LB008715

Walgreens Pharmacist Recommended.
Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
*This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademarks Mucinex® and Fast-Max®.

Drug Facts

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Diphenhydramine HCl 25 mg.....	Antihistamine/cough suppressant
Phenylephrine HCl 10 mg.....	Nasal decongestant

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

Uses ■ temporarily relieves these common cold and flu symptoms
 ■ cough ■ nasal congestion
 ■ minor aches and pains ■ sore throat
 ■ headache ■ runny nose ■ sneezing
 ■ temporarily reduces fever
 ■ controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
 ■ with any other drug containing acetaminophen

Drug Facts (continued)

(prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ with any other drug containing diphenhydramine, even one used on the skin
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 ■ for children under 12 years of age
Ask a doctor before use if you have
 ■ liver disease ■ heart disease
 ■ high blood pressure ■ thyroid disease
 ■ diabetes ■ glaucoma
 ■ trouble urinating due to an enlarged prostate gland
 ■ a breathing problem such as emphysema or chronic bronchitis
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
 ■ cough that occurs with too much phlegm (mucus)
Ask a doctor or pharmacist before use if
 ■ you are taking the blood thinning drug warfarin
 ■ you are taking sedatives or tranquilizers

Drug Facts (continued)

When using this product
 ■ do not use more than directed
 ■ excitability may occur, especially in children
 ■ marked drowsiness may occur
 ■ avoid alcoholic drinks
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
 ■ be careful when driving a motor vehicle or operating machinery
Stop use and ask a doctor if
 ■ nervousness, dizziness, or sleeplessness occur
 ■ pain, nasal congestion, or cough gets worse, or lasts more than 7 days
 ■ fever gets worse, or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur
 ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions
 ■ do not take more than directed (see Overdose warning)
 ■ do not take more than 6 doses in any 24-hour period
 ■ measure only with dosing cup provided. Do not use any other dosing device.
 ■ mL = milliliter ■ keep dosing cup with product
 ■ dose as follows or as directed by a doctor
 ■ adults and children 12 years and older:
 20 mL every 4 hours while symptoms last
 ■ children under 12 years of age: do not use
Other information
 ■ each 20 mL contains: sodium 12 mg
 ■ store between 20-25°C (68-77°F). Do not refrigerate.
Inactive ingredients citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum
Questions or comments?
 Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

PEEL CORNER FOR MORE DRUG FACTS

Print/Varnish Omit Area

DAY & NIGHT PACK		NDC 0363-8994-48
		
DAYTIME • NON-DROWSY	Compare to Vicks® DayQuil™ & NyQuil™ Severe Cold & Flu Relief LiquiCaps™ active ingredients††	
Severe Cold & Flu	NIGHTTIME	NEW
Severe Cold & Flu	Severe Cold & Flu	
ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT GUAIFENESIN 200 mg / EXPECTORANT PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT	ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT DOXYLAMINE SUCCINATE 6.25 mg / ANTIHISTAMINE PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT	
MAXIMUM STRENGTH	MAXIMUM STRENGTH	
ACTUAL SIZE 	ACTUAL SIZE 	
32 SOFTGELS* (*LIQUID-FILLED CAPSULES)	48 TOTAL SOFTGELS	16 SOFTGELS* (*LIQUID-FILLED CAPSULES)

DAY & NIGHT PACK			NDC 0363-8995-24
		<i>Compare to Vicks® DayQuil® Severe Cold & Flu & Vicks® NyQuil® Severe Cold & Flu active ingredients††</i>	
DAYTIME • NON-DROWSY		NIGHTTIME	NEW
Severe Cold & Flu		Severe Cold & Flu	
ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT GUAIFENESIN / EXPECTORANT PHENYLEPHRINE HCl / NASAL DECONGESTANT		ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT DOXYLAMINE SUCCINATE / ANTIHISTAMINE PHENYLEPHRINE HCl / NASAL DECONGESTANT	
✔ DISSOLVES QUICKLY			
MAXIMUM STRENGTH		MAXIMUM STRENGTH	
			
ACTUAL SIZE		ACTUAL SIZE	
16	SOFTGELS	24	SOFTGELS
		8	SOFTGELS

3257

NDC 0363-4150-08

Drug Facts**Active ingredients (in each 15 mL)**

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Purposes**Uses**

■ temporarily relieves common cold/flu symptoms ■ nasal congestion ■ sinus congestion and pressure ■ minor aches and pains ■ headache ■ fever ■ sore throat ■ reduces swelling of nasal passages ■ cough due to minor throat and bronchial irritation ■ temporarily restores freer breathing through the nose ■ promotes nasal and/or sinus drainage ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if: ■ adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount ■ child takes more than 4 doses (15 mL each) in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

← PEEL HERE **Drug Facts** (continued under label)

Walgreens

Compare to the active ingredients in
Vicks® DayQuil® Severe Cold & Flu^{††}

DAYTIME • NON-DROWSY**Severe Cold & Flu**

ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT
GUAIFENESIN 200 mg / EXPECTORANT
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

Maximum Strength

- Relieves headache, fever, sore throat, minor aches & pains; nasal/sinus congestion & sinus pressure; cough & chest congestion
- For ages 6 years & over
- Alcohol free

8 FL OZ (237 mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

PARENTS:
Learn about your medicine above
www.StopMedicineAbuse.org

Discard Seal,
Empty &
Replace Cap



Walgreens Company

[†]Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
^{††}This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

DISTRIBUTED BY: **WALGREEN CO.**
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2022 Walgreen Co.

ITEM 271599 W10064-0922-F



PLD-8340B LB008969

W3ORG0922-F

Drug Facts (continued)

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if the user has ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ a sodium-restricted diet

Ask a doctor or pharmacist before use if user is taking the blood thinning drug warfarin.

When using this product, do not take more than directed.

Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur ■ pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ new symptoms occur ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ take only as directed (see **Overdose warning**) ■ do not take more than 4 doses in any 24-hour period ■ measure only with dosing cup provided. Do not use any other dosing device. ■ mL = milliliter ■ keep dosing cup with product

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information ■ each 15 mL contains: sodium 12 mg ■ store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Walgreens

Compare to Maximum Strength Mucinex® SINUS-MAX®
Severe Congestion Relief active ingredients††

NDC 0363-9018-16

NEW

Severe Congestion Relief

ACETAMINOPHEN 325 mg / PAIN RELIEVER
DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT
GUAIFENESIN 200 mg / EXPECTORANT
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

MAXIMUM STRENGTH

- Relieves headache, sinus pressure & congestion
- Thins & loosens mucus
- Controls cough
- 12 years & older

16 SOFTGELS*
(*LIQUID-FILLED CAPSULES)



ACTUAL SIZE

Walgreens

DAYTIME
NON-DROWSY

Compare to Vicks®
DayQuil® Severe Cold & Flu
active ingredients††

NDC 0363-8996-24

Severe Cold & Flu

ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT
GUAIFENESIN 200 mg / EXPECTORANT
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

✓ **DISSOLVES QUICKLY**

MAXIMUM STRENGTH

- Relieves headache, fever, sore throat, minor aches & pains, cough, chest congestion, nasal/sinus congestion & sinus pressure

24 SOFTGELS



Walgreens

NIGHTTIME

Severe Cold & Flu

Compare to Vicks®
NyQuil® Severe Cold & Flu
active ingredients††

NDC 0363-8997-24

ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT
DOXYLAMINE SUCCINATE 6.25 mg / ANTIHISTAMINE
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

✓ **DISSOLVES QUICKLY**

MAXIMUM STRENGTH

- Relieves headache, fever, sore throat, minor aches & pains, nasal/sinus congestion & sinus pressure, sneezing, runny nose & cough

24 SOFTGELS



DAY & NIGHT PACK

Walgreens

DAYTIME • NON-DROWSY

Severe Sinus

CONGESTION & COUGH

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

• Pain reliever, fever reducer, cough suppressant & nasal decongestant

12 SOFTGELS

Walgreens

NIGHTTIME

Severe Sinus

CONGESTION & COUGH

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
DOXYLAMINE SUCCINATE / ANTIHISTAMINE
PHENYLEPHRINE HCl / NASAL DECONGESTANT

• Relieves sinus congestion & pressure, headache, runny nose, sneezing & cough

8 SOFTGELS

DO NOT TAKE THESE PRODUCTS AT THE SAME TIME

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

DO NOT TAKE THESE PRODUCTS AT THE SAME TIME

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS


Walgreens

SEVERE SINUS CONGESTION & COUGH DAYTIME SOFTGELS

Walgreens

SEVERE SINUS CONGESTION & COUGH NIGHTTIME SOFTGELS

Walgreens



Walgreens

Compare to Maximum Strength Mucinex® Sinus-Max® Severe Congestion Relief active ingredients^{††}

NDC 0363-0524-16

Severe Sinus Congestion

ACETAMINOPHEN 325 mg / PAIN RELIEVER
DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT
GUAIFENESIN 200 mg / EXPECTORANT
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

MAXIMUM STRENGTH

- Relieves sinus congestion & headache
- Thins & loosens mucus
- Controls cough
- Ages 12 & older
- Alcohol free

16 SOFTGELS*
 (*LIQUID-FILLED CAPSULES)



ACTUAL SIZE

DISTRIBUTED BY: **WALGREEN CO.**
 200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
 walgreens.com
 ©2017 Walgreen Co.

ITEM 984982 W10024-0118-L



FD-3426B
 F000469

Lot No.:
 Exp. Date:

Drug Facts

Active ingredients (in each softgel)	Purposes
Acetaminophen 325 mg	Pain reliever
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves
- cough due to inhaled irritants
- nasal congestion
- sinus congestion and pressure
- headache
- minor aches and pains
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash. If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- diabetes
- high blood pressure
- heart disease
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Drug Facts (continued)

■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 softgels in any 24-hour period
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use

Other information

- swallow whole, do not crush, chew, or dissolve
- store between 15°-30°C (59°-86°F)
- avoid excessive heat

Inactive ingredients FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, titanium dioxide

Questions or comments?
 Call 1-877-753-3935 Monday-Friday 9AM-5PM EST



Health expertise
 you rely on,
 quality you trust.

[†]Walgreens Pharmacist Survey
^{††}This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Sinus-Max® Severe Congestion Relief.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Walgreens

Compare to Maximum Strength
Mucinex® Sinus-Max® Severe
Congestion Relief active ingredients^{††}

SINUS RELIEF NDC 0363-0478-06

**Severe
Congestion**

ACETAMINOPHEN 650 mg / PAIN RELIEVER
GUAIFENESIN 400 mg / EXPECTORANT
PHENYLEPHRINE HCl 10 mg /
NASAL DECONGESTANT

MAXIMUM STRENGTH

- Relieves headache & sinus congestion
- Thins & loosens mucus
- 12 years & older

PLD-B388A
LB004975

6 FL OZ (177 mL) LIQUID

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING

Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey

^{††}This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Sinus-Max® Severe Congestion Relief.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2017 Walgreen Co.

ITEM 923791 W10013-0917-L

3 11917 18102 8

PLD-B388A
LB004976

ORG0817-F

Drug Facts

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg.....	Pain reliever
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

Uses

- temporarily relieves
 - headache ■ nasal congestion
 - minor aches and pains
 - sinus congestion and pressure
 - temporarily promotes nasal and/or sinus drainage
 - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease ■ heart disease ■ diabetes
- thyroid disease ■ high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product ■ mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years and older: 20 mL every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, EDTA disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

age	dose
children 6 to under 12 years of age	10 mL every 4 hours
children 4 to 5 years	5 mL every 4 hours
children under 4 years	do not use

Other information

- each 5 mL contains: sodium 2 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients acesulfame potassium, alcohol, benzoic acid, citric acid, disodium EDTA, FD&C red #40, flavor, maltitol, propylene glycol, purified water, sodium citrate

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2015 Walgreen Co.

W10040-XXXX-F
ITEM 448737
PLD-F319C LB003474
REV1013-RF_R01



children's
DAYTIME
Cold & Cough
Dextromethorphan HBr /
Cough Suppressant
Phenylephrine HCl /
Nasal Decongestant
• **ALCOHOL 0.14%**




4 FL OZ (118 mL)
CHERRY FLAVOR

NDC 0363-3780-04

Drug Facts

Active Ingredients (in each 5 mL)

Dextromethorphan HBr 5 mg.....Cough suppressant
Phenylephrine HCl 2.5 mg.....Nasal decongestant

Purposes

Uses

- temporarily relieves
- nasal and sinus congestion
- cough due to minor throat and bronchial irritation as may occur with a cold

Warnings

Do not use ■ in a child under 4 years of age ■ in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product. **Ask a doctor before use if the child has** ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ a breathing problem such as chronic bronchitis ■ cough that occurs with too much phlegm (mucus) ■ chronic cough that lasts, or as occurs with asthma. **When using this product, do not use more than directed. Stop use and ask a doctor if** ■ nervousness, dizziness or sleeplessness occur

- symptoms do not improve within 7 days or occur with fever
- cough persists more than 7 days, comes back or occurs with a fever, rash or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- shake well before using

TAMPER EVIDENT: DO NOT USE IF PRINTED BURST SEAL UNDER CAP IS TORN OR MISSING

Drug Facts

Active Ingredients (in each 20 mL) **Purposes**
 Acetaminophen USP 650 mg.....Pain/fever reducer
 Dextromethorphan HBr USP 20 mg.....Cough suppressant
 Guaifenesin USP 400 mg.....Expectorant
 Phenylephrine HCl USP 10 mg.....Nasal decongestant

Uses

- Temporarily relieves these symptoms occurring with a cold or flu:
- cough due to minor throat and bronchial irritation
- nasal congestion
- sore throat and hoarseness
- minor aches and pains
- fever
- temporarily reduces fever
- helps loosen phlegm (mucus) and the bronchial passages to drain
- helps thin mucus and makes coughs more productive

Warnings

- Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take:
 - more than 4 doses in 24-hour period, which is the maximum daily amount
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks every day while using this product
- Always alert your doctor if you are taking any other drugs. Symptoms may include:
 - nausea/vomiting
 - loss of appetite
 - dark urine
 - itching
 - fatigue
 - joint pain
 - stomach pain
 - yellowing of the skin or eyes
 - dark stools
- If a skin rash occurs, stop use and seek medical help right away. Some throat warnings: If your throat is sore, painful for more than 2 days, is accompanied by fever, swelling, or difficulty swallowing, consult a doctor promptly.

Do not use

- If you are now taking a prescription for non-sedative cough medicine (NAC), sedative cough medicine, sedative, or sedative and have or have been diagnosed, or for 2 weeks after stopping the NAC drug. If you do not know if your prescription drug contains an NAC, ask a doctor or pharmacist before taking this product.
- If you are taking any other drug containing acetaminophen or guaifenesin. If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- Ask a doctor before use if you have:
 - liver disease
 - heart disease
 - high blood pressure
 - thyroid disease
 - diabetes
 - trouble urinating
 - due to an enlarged prostate gland
 - cough that occurs with the heart
 - allergy to any of the ingredients
 - if you are pregnant or planning to get pregnant
 - if you are breastfeeding
 - if you are taking any other drug containing acetaminophen or guaifenesin

- Ask a doctor or pharmacist before use if you are:
 - taking the blood thinning drug warfarin
 - taking any other oral nasal decongestant or antihistamine
 - taking any other pain reliever/fever reducer

When using this product do not use more than directed

- Stop use and ask a doctor if:
 - if you get nervous, dizzy, or sleepy
 - if you get a rash or redness, hives, or itching
 - if you get a headache or dizziness
 - if you get a sore throat or difficulty swallowing
 - if you get a fever or chills
 - if you get a cough that does not go away
 - if you get a cough that is worse than when you started
 - if you get a cough that is worse than when you started
 - if you get a cough that is worse than when you started

If pregnant or breast-feeding, ask your health care provider before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Prompt medical attention is critical for your health. Ask your doctor or pharmacist if you have any other symptoms.

Drug Facts (continued)

Directions

- Do not take more than 4 doses in any 24-hour period.
- Do not exceed recommended dosage. Taking more than the recommended dose increases the risk of liver damage.
- Measure only with marked cup provided.
- Keep dosing cup with product.
- Keep out of reach of children under 12 years of age.

Age	Dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

Other information

- Each 20 mL contains acetaminophen 7 mg.
- Contains 10 mg of alcohol per 20 mL.

Inactive ingredients include: citric acid, disodium edetate, FD&C Yellow No. 6, glycerin, hydroxyethylcellulose, polyethylene glycol, propyl paraben, purified water, sodium benzoate, sodium lactate, sucralose, and xanthan gum.

Questions or comments? 1-866-467-2748

IMPORTANT: Keep this carton for future reference on NDI labeling.



Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, owner of the registered trademark Adult Robitussin Maximum Strength SEVERE Multi-Symptom Cough Cold & Flu CF MAX.

DISTRIBUTED BY: WALGREEN CO.
 200 WILNOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED

walgreens.com
 ©2022 Walgreen Co.



ADULT SEVERE

Tussin CF Max

NDC 3383-5306-00



Compare to the active ingredients in Adult Robitussin Maximum Strength SEVERE Multi-Symptom Cough Cold & Flu CF MAX.

ADULT SEVERE

Tussin CF Max

MULTI-SYMPTOM COUGH, COLD & FLU

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
 DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
 GUAFENESIN / EXPECTORANT
 PHENYLEPHRINE HCl / NASAL DECONGESTANT

Maximum Strength

Sugar Free

Alcohol Free

- Relieves cough, sore throat, body aches, fever, nasal congestion & chest congestion
- 12 years & older



8 FL OZ (237 mL) Natural Raspberry flavor

63008WAGFC

Lot:

Exp:

ITEM 546870 W00000-0000-0



3 11917 12384 4



Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back, or occurs with rash or headache that lasts.
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • take only as directed

• only use the dose cup provided • do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
---------------------------------	-------------------

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information:

Inactive ingredients: citric acid, D&C Yellow No.10, disodium edetate, FD&C Blue No. 1, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose

Questions or comments? 1-866-467-2748

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® Dayquil™ Severe+ VapoCOOL™.

LOT: 72712WAGLR

DIST. BY: WALGREEN CO. 200 WILMOT RD.
DEERFIELD, IL 60015 100% SATISFACTION
GUARANTEED walgreens.com ©2021 Walgreen Co.
ITEM 355983 W00000-0000-0

EXP:

3 ■■■ 11917 ■ 03128 ■ ■ 6

NDC 0363-7270-12

Walgreens

Compare to the active ingredients in
Vicks® DayQuil™ Severe+ VapoCOOL™¹

DAYTIME

Severe Cold & Flu

ACETAMINOPHEN / MINOR ACHES & PAINS / FEVER
PHENYLEPHRINE HCl / NASAL CONGESTION /
SINUS PRESSURE
DEXTROMETHORPHAN HBr / COUGH
GUAIFENESIN / CHEST CONGESTION

- Pain reliever, fever reducer, cough suppressant, expectorant, nasal decongestant

12 FL OZ (354 mL)

Drug Facts

Tamper Evident: Do not use if imprinted shrink band is missing or broken

Active ingredients (in each 30 mL)

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms: • nasal congestion • sinus congestion & pressure • cough due to minor throat & bronchial irritation • minor aches & pains • headache • fever • sore throat • reduces swelling of nasal passages • temporarily restores freer breathing through the nose • promotes nasal and/or sinus drainage • helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 4 doses in 24 hrs, which is the maximum daily amount for this product • taken with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product.

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
• Skin reddening • Blisters • Rash

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema • a sodium-restricted diet ►

Drug Facts (continued)

When using this product • do not use more than directed • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks

• be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • take only as directed

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information • each 30 mL contains: sodium 81 mg • store at room temperature. Do not refrigerate.

Inactive ingredients citric acid, D&C Yellow No. 10, disodium edetate, FD&C Blue No. 1, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose.

Questions or comments? 1-866-467-2748

†Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
‡This product is not manufactured or distributed by Doctor & Gamble, distributor of Vicks® NyQuil™ Severe® VapoCOOL™.

LOT: 61412WAGLR

DIST. BY: WALGREEN CO. 200 WILMOT RD.,
DEERFIELD, IL 60015 100% SATISFACTION
GUARANTEED walgreens.com ©2021 Walgreen Co.
ITEM 355983 W00000-0000-0

EXP: 3 11917 03128 6

NDC 0363-6140-12

Walgreens

Compare to the active ingredients in
Vicks® NyQuil™ Severe® VapoCOOL™

NIGHTTIME

**Severe
Cold & Flu**

ACETAMINOPHEN / MINOR ACHES & PAINS / FEVER
PHENYLEPHRINE HCl / NASAL CONGESTION /
SINUS PRESSURE
DOXYLAMINE SUCCINATE / SNEEZING / RUNNY NOSE
DEXTROMETHORPHAN HBr / COUGH

Severe Cooling

• Pain reliever, fever reducer,
nasal decongestant,
antihistamine,
cough suppressant

12 FL OZ (354 mL)

Drug Facts **DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN.**

Active ingredients (in each 30 mL)

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine
Phenylephrine HCl 10 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms: • nasal congestion • sinus congestion & pressure • cough due to minor throat & bronchial irritation • cough to help you sleep • minor aches & pains • headache • fever • sore throat • runny nose & sneezing • reduces swelling of nasal passages • temporarily restores freer breathing through the nose • promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 4 doses in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • Skin reddening • Blisters • Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema • trouble urinating due to enlarged prostate gland • a sodium-restricted diet

Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers • taking the blood thinning drug warfarin.



Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults) • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

• take only as directed

• only use the dose cup provided • do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over 30 mL every 4 hrs

children 6 to under 12 yrs 15 mL every 4 hrs

children 4 to under 6 yrs ask a doctor

children under 4 yrs do not use

Other information • each 15 mL contains: sodium 9 mg • store at room temperature. Do not refrigerate.

Inactive ingredients citric acid, D&C Yellow No.10, edetate disodium, FD&C Green No. 3, FD&C Red No. 40, flavors, glycerin, menthol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum.

Questions or comments? 1-866-467-2748

*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® Dayquil™ Severe Cold & Flu Honey Flavor.

LOT: 61012XXLR Distributed by:

UNVARNISHED
AREA

EXP:



*Compare to the active ingredients in
Vicks® Dayquil™ Severe Cold & Flu
Honey Flavor
NDC 0363-6110-12

DayTime Cold & Flu Relief

Acetaminophen - Pain reliever/Fever reducer
Guaifenesin - Expectorant
Phenylephrine HCl - Nasal decongestant
Dextromethorphan HBr - Cough suppressant

- Headache, Fever, Sore Throat, Minor Aches & Pains
- Chest Congestion, Thins & Loosens Mucus
- Nasal Congestion, Sinus Pressure
- Cough

Honey Flavor

Naturally and Artificially Flavored

12 FL OZ (354 ml)

Drug Facts

DO NOT USE IF IMPRINTED SHRINK BAND IS
MISSING OR BROKEN

Active ingredients (in each 15 mL)

	Purpose
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms: • nasal congestion • sinus congestion & pressure • cough due to minor throat & bronchial irritation • minor aches & pains • headache • fever • sore throat • reduces swelling of nasal passages • temporarily restores freer breathing through the nose • promotes nasal and/or sinus drainage • helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if • adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product • child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product • taken with other drugs containing acetaminophen • adult has 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • Skin reddening • Blisters • Rash

If a skin reaction occurs, stop use and seek medical help right away
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

TAMPER EVIDENT: Do not use if printed shrink band is missing or broken

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 100 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves common cold symptoms:
 - nasal congestion
 - sinus congestion & pressure
 - cough due to minor throat & bronchial irritation
 - cough to help you sleep
 - reduces swelling of nasal passages
 - temporarily restores freer breathing through the nose
 - promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease

This product is not manufactured or distributed by Procter & Gamble, the distributor of Vicks® DayQuil™ Kids Honey Cold & Cough + Mucus.

Walgreens

Compare to the active ingredients in Vicks® DayQuil™ Kids Honey Cold & Cough + Mucus

DAYTIME

Children's Cold, Cough & Mucus

GUAIFENESIN / EXPECTORANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT

Acetaminophen Free

+ Honey

Chest congestion, nasal congestion & cough

Honey naturally & artificially flavored

8 FL OZ (236 mL)

Drug Facts (continued)

- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema

When using this product, do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not improve within 7 days or occur with a fever
- cough persists for more than 7 days, comes back or occurs with a fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

Walgreens Pharmacist Recommended. Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2023 Walgreen Co. 62308WAGLR

PLASTIC BOTTLE PLASTIC CUP

W30H0423-F

ITEM 206909 W00000-0000-0

LOT: UNVARNISHED AREA

EXP:

3 11917 00697 0

PEEL BACK HERE FOR DRUG FACTS

Drug Facts (continued)

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs




adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 12 mg
- store at room temperature

Inactive ingredients anhydrous citric acid, carboxymethyl cellulose sodium, D&C Yellow No. 10, disodium edetate, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum.

Questions or comments? 1-866-467-2748



6.75" Width

3.25" Height

6.75" Width

Questions or comments? 1-800-467-2748

READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE. KEEP CARTON FOR REFERENCE. DO NOT DISCARD.

Drug Facts

Active ingredients (in each packet)	Purpose
Acetaminophen 500 mg	Pain reliever/fever reducer
Dextromethorphan hydrobromide 20 mg	Cough suppressant
Phenylephrine hydrochloride 10 mg	Nasal decongestant

Uses

- Temporarily relieves these symptoms due to a cold:
 - minor aches and pains
 - headache
 - minor sore throat pain
 - nasal and sinus congestion
 - cough due to minor throat and bronchial irritation
- Temporarily reduces fever

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 packets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include skin redness, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult physician promptly.

Do not use

- in a child under 4 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription and nonprescription). Ask a doctor or pharmacist before using with other drugs if you are not sure.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product

- do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- nausea or swelling is present
- new symptoms occur
- symptoms do not get better or worsen
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

CONVENIENT RECLOSING TAB

MULTI-SYMPTOM
Wal-Flu®
Severe Cold

Well at
Walgreens
WALGREENS PHARMACIST RECOMMENDED*

NEW Well at
Walgreens

WALGREENS PHARMACIST RECOMMENDED*

MULTI-SYMPTOM
Wal-Flu®
Severe Cold
Acetaminophen /
Pain Reliever / Fever Reducer
Dextromethorphan HBr /
Cough Suppressant
Phenylephrine HCl /
Nasal Decongestant

Relieves:

- Nasal congestion
- Cough
- Body ache
- Sore throat pain
- Headache
- Fever


GREEN TEA & HONEY LEMON FLAVORS

Compare to Theraflu® Multi-Symptom Severe Cold active ingredients^{1†}

6 PACKETS

*Walgreens Pharmacist Survey Study, November 2014.
†This product is not manufactured or distributed by Novartis, owner of the registered trademark Theraflu®.

Questions or comments?
1-800-925-4733

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED

walgreens.com ©2016 Walgreen Co.
MADE IN CANADA

ITEM 0003016
11917177540

C-3192-30508-BW-1

ORG0515-F2

Walgreens

Compare to Vicks® DayQuil®
LiquiCaps® active ingredients!!

NDC 0363-8998-24

DAYTIME • NON-DROWSY

Cold & Flu

ACETAMINOPHEN 325 mg / ACHES / FEVER / SORE THROAT
DEXTROMETHORPHAN HBr 10 mg / COUGH
PHENYLEPHRINE HCl 5 mg / NASAL CONGESTION

MULTI-SYMPTOM

LIQUID CAPS

- Relieves aches, fever, sore throat,
nasal congestion & cough

24

LIQUID CAPS



ACTUAL SIZE

3277